



Does early exposure to a visual signed language “hurt” auditory language tissue development: Evidence from fNIRS neuroimaging of language processing in deaf individuals with cochlear implants

Kajia K. Jasinska, Clifton Langdon, and Laura-Ann Pettito

Pettito Brain & Language Laboratory (BL²) and Visual Language and Visual Learning, VL2, NSF Science of Learning Center

NSF Center Funding: SBE 1041725

Funding: Pettito (P) NIH SR01HD45822 and NIH SR21HD050558, Canadian Foundation for Innovation, Ontario Research Fund

ABSTRACT

Controversy abounds regarding the specific impact of differences in language experience on the acquisition of spoken language in deaf individuals with cochlear implants (CI)^{1,2,3,4}. Noteworthy are claims that due to neuroplasticity of auditory language tissue, early exposure to a signed language results in deviance to auditory language tissue development. We find that early, but not later, exposure to a signed language supports typical language development.

QUESTION

Does early exposure to a visual signed language impact classic left-hemisphere spoken language tissue development, including left Inferior Frontal Gyrus (LIFG) and Superior Temporal Gyrus (STG)^{5,6,7}, in a deaf individual with a CI?

Does age of signed language exposure impact language processing in the deaf CI individual?

HYPOTHESES

- H1 Early language exposure, *both signed and spoken*, facilitates normal neural development for language processing
- P1 CI individual with *early, but not later*, signed language exposure recruit left-hemisphere language areas
- H2 *Only* early spoken language exposure facilitates normal neural development for language processing
- Early signed language exposure disrupts development
- P2 CI individual with *early and later* signed language exposure do *not* recruit left-hemisphere language areas

METHODS

Participants

Deaf Adults with cochlear implant
(Age of implant = 13 years)




Early Signed Language Exposure
(ASL before age 5 years)
N = 4

Later Signed Language Exposure
(ASL after age 5 years)
N = 4

Task


Reading aloud English words
Near Infrared Spectroscopy
(fNIRS) neuroimaging^{8,9}

Advancement: Optical imaging
is safe for cochlear implant
users




RESULTS

Early Signed Language Exposed Deaf CI Individuals Show Greater Neural Activation in Left Inferior Frontal Gyrus



Later Signed Language Exposed Deaf CI Individuals Show Greater Neural Activation in Right Inferior Frontal Gyrus



CONCLUSION

Early exposed deaf CI individuals showed greater activation in classic **left-hemisphere** language areas (LIFG)

Late exposed deaf CI individuals showed greater activation in the **right-hemisphere** (RIFG), not in classic left hemisphere language areas (LIFG)

Supports Hypothesis 1

Early signed language exposure facilitates normal language processing

No evidence of a negative impact on language processing as a result of early visual signed language exposure

Implications

Optimal developmental timing of signed language exposure

Early language exposure, be it signed or spoken, supports healthy, typical language development

New view on how early life language exposure, irrespective of modality (signed, spoken), can facilitate language processing in the deaf CI individual

REFERENCES

- Seal, B. C., Nussbaum, D. B., Betzner, K. A., Scott, S., & Waddy-Smith, B. (2011). Cochlear implants international. 12(1), 34–43.
- Osberger, M. J., Miyamoto, R. T., Zimmerman-Phillips, S., Kemink, J. L., Stree, B. S., Firszt, J. S., & Novak, M. A. (1997). Ear and Hearing, 12(4), 685–805.
- Svirsky, M. A., Robbins A.M., Kirk K.L., Pisoni D.B., Miyamoto R.T. (2000). Psychological Science, 11(2), 153–158.
- Visual Language and Visual Learning Science of Learning Center. (2012, June). Washington, DC: J. Mitchner, D.B. Nussbaum, and S. Scott.
- Pettito, Berens, Koveinman, et al. (2012). Brain & Lang.
- Pettito, Zatorre, et al. (2000). Proc. Natl. Acad. Sci. 97, 13961–13966
- Koveinman, Baker, & Pettito (2008). J. Cog. Neurosci. 20, 153–169
- Shalinsky, Koveinman, Berens, & Pettito (2010) J. Vis. Exp. 29. http://www.jove.com/index/details.asp?id=1268
- Ye, Tak, Jiang, Jung, & Jang (2009). Neuroimaging, 44, 428–447
- Woodcock (1991). Woodcock Language Proficiency Battery - Revised

*CORRESPONDING AUTHOR

Laura-Ann Pettito@gallaudet.edu; www.pettito.gallaudet.edu
Visual Language & Visual Learning, VL2; and Brain and Language Laboratory, BL2. Society for Neuroscience, November 9-13, 2013